

Supporting Information

The Effect of Hydrofluoroether Cosolvent Addition on Li Solvation in Acetonitrile-Based Solvate Electrolytes and Its Influence on S Reduction in a Li-S battery

Kimberly A. See,^{#,±,†} Heng-Liang Wu,^{#,±,†} Kah Chun Lau,^{±,†} Minjeong Shin,^{±,†} Lei Cheng,^{±,‡} Mahalingam Balasubramanian,^{±,∞} Kevin G. Gallagher,^{±,§} Larry A. Curtiss,^{±,‡} and Andrew A. Gewirth^{*,±,†}

[#] These authors contributed equally to this work.

[±] Joint Center for Energy Storage Research, 9700 S. Cass Avenue, Argonne, IL 60439

[†] Department of Chemistry, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801 United States

[‡] Department of Physics and Astronomy, California State University, Northridge, California 91330 United States

[§] Materials Science Division; [§] Chemical Sciences and Engineering Division; [∞] X-ray Science Division, Advanced Photon Source; Argonne National Laboratory, Argonne, Illinois 60439 United States

Corresponding Author:

*Email: agewirth@illinois.edu




Figure S1. (a) ^7Li and (b) ^{19}F NMR of wet $\text{MeCN}_{2.5}\text{-LiTFSI}$ (>50 ppm water) prepared with as-purchased LiTFSI before and after exposure to Li metal. A significant shift in the resonance of both ^7Li and ^{19}F is observed after exposure to Li metal due to the decomposition of the electrolyte. A visible color change is observed as shown by the photograph inset in (a). The dry $\text{MeCN}_{2.5}\text{-LiTFSI}$ (<15 ppm water) prepared with dried LiTFSI does not exhibit a color change upon exposure to Li metal.




Figure S2. The first 10 cyclic voltammograms (CV) of a Li-S@CMK-3 cells scanned at 0.1 mV s^{-1} with (a) the solvate electrolyte, $(\text{MeCN})_2\text{-LiTFSI}$, and the solvate diluted with TTE at volume ratios of (b) 2:1, (c) 1:1, and (d) 1:2 (solvate:TTE).




Figure S3. Raman spectra of several concentrations of LiTFSI in MeCN up to the $(\text{MeCN})_2\text{-LiTFSI}$ at 3.98 M. The concentrations shown are 3.70 M, 2.47 M, 1.85 M, 0.93 M, and 0.46 M. (a) The lower wavenumber region reveals a shift and intensity increase in the TFSI mode, a, to a new mode, mode b, due to coordinated TFSI. (b) The higher wavenumber region shows MeCN modes c, d, e, and f. Modes d and f grow in at the expense of mode c and e, respectively, due to MeCN coordination to Li^+ . Mode assignments can be found in Table S1.

Table S1. Assignments of Raman modes in the LiTFSI concentration series in MeCN

ID	Raman shift (cm^{-1})							Species	Assignment	Ref.
	LiTFSI concentration in MeCN (M)									
	0.46	0.93	1.85	2.47	3.08	3.70	3.98			
a	739	739	740	739	739	739	739	TFSI	S-N stretch + C-S stretch + CF_3 bend	1-4
b	<i>w</i>	<i>w</i>	<i>w</i>	742	744	746	746	TFSI	mode a, coordinated to Li^+	1-4
c	2253	2251	2252	2253	2253	2256	2253	MeCN	$\text{C}\equiv\text{N}$ stretch	5,6
d	2276	2276	2276	2276	2277	2278	2278	MeCN	mode c, coordinated to Li^+	7-9
e	2291	2291	2293	2293	2293	2294	2296	MeCN	C-H stretch	5,6
f	<i>i</i>	2305	2308	2306	2307	2308	2308	MeCN	mode e, coordinated to Li^+	

w indicates the mode was either absent or too weak to determine the position.




Figure S4. Raman spectra of the (a) $(\text{MeCN})_{2.5}\text{-LiTFSI}$ and (b) $(\text{MeCN})_2\text{-LiTFSI}$ neat and diluted with TTE at volume ratios of 2:1, 1:1, and 1:2 (electrolyte:TTE). The spectra are fit with four component Gaussian functions to determine the relative peak area ratios associated with free MeCN and coordinated MeCN.




Figure S5. Relative ${}^7\text{Li}$ chemical shifts calculated by DFT for LiTFSI in MeCN in the dilute regime, the $(\text{MeCN})_2\text{-LiTFSI}$ solvate, and $(\text{MeCN})_2\text{-LiTFSI}$ diluted with TTE at a volume ratio of 1:1. As the LiTFSI concentration increases from the dilute regime to the $(\text{MeCN})_2\text{-LiTFSI}$ solvate, the ${}^7\text{Li}$ chemical shift becomes more positive (deshields). Addition of TTE to the $(\text{MeCN})_2\text{-LiTFSI}$ solvate also causes the ${}^7\text{Li}$ chemical shift to move to a more deshielded region. The shift to more positive ppm as the LiTFSI concentration increases and TTE is added is consistent with the experimental results.

References

- (1) Bakker, A.; Gejji, S.; Lindgren, J.; Hermansson, K.; Probst, M. M. *Polymer* **1995**, *36*, 4371–4378.
- (2) Edman, L. *J. Phys. Chem. B* **2000**, *104*, 7254–7258.
- (3) Herstedt, M.; Smirnov, M.; Johansson, P.; Chami, M.; Grondin, J.; Servant, L.; Lassègues, J. C. *J. Raman Spectrosc.* **2005**, *36*, 762–770.
- (4) Brouillette, D.; Irish, D. E.; Taylor, N. J.; Perron, G.; Odziemkowski, M.; Desnoyers, J. E. *Phys. Chem. Chem. Phys.* **2002**, *4*, 6063–6071.
- (5) Neelakantan, P. *Proc. Indian Acad. Sci. - Sect. A* **1964**, *60*, 422–424.
- (6) Fini, G.; Mirone, P. *Spectrochim. Acta Part Mol. Spectrosc.* **1976**, *32*, 439–440.
- (7) Han, S.-D.; Borodin, O.; Seo, D. M.; Zhou, Z.-B.; Henderson, W. A. *J. Electrochem. Soc.* **2014**, *161*, A2042–A2053.
- (8) Barthel, J.; Buchner, R.; Wismeth, E. *J. Solut. Chem.* **29**, 937–954.
- (9) Brouillette, D.; Irish, D. E.; Taylor, N. J.; Perron, G.; Odziemkowski, M.; Desnoyers, J. E. *Phys. Chem. Chem. Phys.* **2002**, *4*, 6063–6071.